

Synthetic LXXLL Peptides Antagonize 1,25-Dihydroxyvitamin D₃-Dependent Transcription

J. Wesley Pike,^{1*} Peterman Pathrose,² Olga Barmina,¹ Ching-Yi Chang,³ Donald P. McDonnell,³ Hironori Yamamoto,¹ and Nirupama K. Shevde¹

¹Department of Biochemistry, University of Wisconsin, Madison, Wisconsin 53706

²Department of Molecular and Cellular Physiology, University of Cincinnati, Cincinnati, Ohio 45267

³Department of Pharmacology and Cancer Biology, Duke University, Durham, North Carolina 27710

Abstract The vitamin D receptor (VDR) is known to mediate the biological actions of 1,25-dihydroxyvitamin D₃ (1,25(OH)₂D₃) through its ability to regulate cellular programs of gene expression. We identified VDR- and retinoid X receptor (RXR)-interacting LXXLL peptides using a mammalian two-hybrid system and examined whether these molecules could block vitamin D and 9-*cis* retinoic acid (9-*cis* RA) response. Peptides were identified that were reactive to RXR alone as well as to both VDR and RXR. Peptide fusion proteins were then examined in MC3T3 E1 cells for their ability to block induction of the osteocalcin promoter by 1,25(OH)₂D₃ or stimulation of an RARE-TK reporter by 9-*cis* RA. Peptides that interacted with both VDR and RXR blocked 1,25(OH)₂D₃-dependent transcription by up to 75%. Peptides that interacted with RXR blocked 9-*cis* RA induced transcription. Two RXR-interacting peptides, however, were also found to block 1,25(OH)₂D₃ response effectively. These studies support the idea that comodulator recruitment is essential for VDR- and RXR-mediated gene expression and that RXR is required for 1,25(OH)₂D₃-induced osteocalcin gene transcription. This approach may represent a novel means of assessing the contribution of RXR in various endogenous biological responses to 1,25(OH)₂D₃. J. Cell. Biochem. 88: 252–258, 2003. © 2002 Wiley-Liss, Inc.

Key words: peptides; VDR; RXR; comodulators; transcription

The vitamin D receptor (VDR) is a member of the steroid and thyroid hormone receptor superfamily of transcription factors that mediates the biological actions of the calcitropic hormone 1,25-dihydroxyvitamin D₃ (1,25(OH)₂D₃) [Evans, 1989; Haussler et al., 1998]. Fundamental advances in our understanding of the mechanism of vitamin D action include the initial cloning and structural characterization of the VDR [Baker et al., 1988; Burmester et al., 1988] definition and characterization of DNA binding sites for the VDR on the OC and OP

promoters [Kerner et al., 1989; Noda et al., 1990], and the discovery that VDR requires a partner protein subsequently identified as retinoid X receptor (RXR) [Sone et al., 1991b; Yu et al., 1991]. The central importance of VDR in calcium and phosphorus homeostasis has been highlighted in the human syndrome of hereditary 1,25(OH)₂D₃-resistant rickets [Malloy et al., 1999].

Active VDR is believed to be composed of a VDR/RXR heterodimer that binds to specific DNA sequences in response to 1,25(OH)₂D₃ [Mangelsdorf et al., 1995]. Early biochemical as well as more recent molecular studies have revealed that the interaction between the VDR and RXR occurs via several α helical regions and is modulated by ligand much like that observed for other nuclear receptors [Whitfield et al., 1995; Jin et al., 1996]. The interaction between RXR and VDR is not dependent upon the presence of nucleic acids, suggesting that ligand-induced formation of the protein complex may precede its interaction with DNA [Sone et al., 1991a; Cheskis and Freedman, 1996; Love et al., 2002].

Grant sponsor: National Institutes of Health; Grant number: DK-52453; Grant sponsor: National Cancer Institute; Grant number: CA 90645; Grant sponsor: Department of Defense; Grant number: DAMD17-99-1-9173.

*Correspondence to: Dr. J. Wesley Pike, Department of Biochemistry, University of Wisconsin, 433 Babcock Drive, Madison, WI 53706. E-mail: pike@biochem.wisc.edu

Received 20 June 2002; Accepted 24 June 2002

DOI 10.1002/jcb.10336

© 2002 Wiley-Liss, Inc.

As with other nuclear receptors, the RXR/VDR heterodimer functions to recruit additional factors that play an essential role in transcription [Rachez et al., 2000]. VDR is able to recruit a group of some 10–13 proteins termed D Receptor Interacting Proteins (DRIP complex) [Rachez et al., 1999; Ryu et al., 1999]. The single protein that mediates interaction with the VDR is DRIP₂₀₅ [Rachez and Freedman, 2000]. VDR is also known to interact with members of the p160 class of coactivators including Grip and Src-1, factors that are important to sex steroid receptor function. These factors appear to act directly to catalyze selective acetylation of histones and indirectly to recruit additional regulatory molecules such as histone acetyltransferases capable of eliciting related changes in chromatin structure [Rachez and Freedman, 2000]. The interaction of the VDR with comodulators such as DRIP₂₀₅ as well as Src-1 and Grip is mediated through LXXLL motifs that associate directly within the AF-2 region of nuclear receptors [McInerney et al., 1998; Perissi et al., 1999]. Recent crystallographic studies have defined the interaction between the LXXLL motif and the AF-2 cleft of the nuclear receptors created upon ligand-binding [Brzozowski et al., 1997; Shiau et al., 1998; Gampe et al., 2000].

Small LXXLL peptides have been identified that target the AF-2 domain of the nuclear receptor family [Chang et al., 1999; Hall et al., 2000]. These peptides mimic the LXXLL motifs found in at least three classes of comodulator proteins, display a certain degree of selectivity for members of the nuclear receptors family and are able to block ligand-induced receptor transactivation in reporter gene assays when linked to a large protein sequence such as the Gal4 DNA binding domain (DBD). We screened a subset of previously identified LXXLL peptides for reactivity with both VDR and RXR and show that several can block vitamin D response. Our results provide additional evidence that the AF-2 region of VDR is essential for 1,25(OH)₂D₃ stimulated transcription and that RXR is a participant in vitamin D response.

MATERIALS AND METHODS

Reagents

Modified Eagles medium alpha (α -MEM) and Dulbecco's modified essential medium (DMEM) were purchased from Life Technologies (Grand Island, NY).

DNA Plasmids

pAVhVDR, pCMX-RXR α , pVDR-VP16, and pRXR α -VP16 were described earlier [Noda et al., 1990; Hall et al., 2000]. phOC 3900 and pGal4(5x)luc3 have been previously reported [Noda et al., 1990; Hall et al., 2000]. pTK-RXRE was a gift from Dr. David Mangelsdorf (University of Texas-Southwestern). pM-Src-1 (residues 621–765), pM-Grip (residues 629–760), pM-C33, pM-D47, pM-EBIP41, pM-EBIP44, pM-EBIP45, pM-EBIP53, pM-EBIP60, pM-EBIP66, pM-EBIP76, pM-EBIP87, pM-EBIP92, and pM-F6 prepared from the pM vector (Clontech, San Diego, CA) containing Gal4 (residues 1–147) were also previously described [Chang et al., 1999; Hall et al., 2000].

Cell Culture

COS7 cells were cultured in DMEM supplemented with 10% fetal bovine serum. MC3T3-E1 cells were cultured in α -MEM supplemented with 10% fetal bovine serum. Transient transfections were carried out in 24-well plates using Lipofectamine Plus as described by the manufacturer. A total of 500–600 ng of DNA per well was introduced into cells. Cells were harvested 24 h later, lysed, and extracts examined for both β -galactosidase and luciferase activities as previously described [Jin et al., 1996]. Luciferase activity was normalized for transfection efficiency using β -galactosidase activity. Individual transfection assays were performed at least three times.

Statistics


Triplicates were averaged and standard deviations determined. Statistical significance was determined using ANOVA.

RESULTS

Src-1 and Grip Peptides Inhibit 1,25(OH)₂D₃ Induced Osteocalcin Promoter Transactivation

We first examined the ability of three tandem cognate LXXLL motifs derived from Src-1 and Grip fused to the larger Gal4 DBD to suppress 1,25(OH)₂D₃ stimulated transactivation in the MC3T3 E1 cell line (Fig. 1a). As can be seen in Figure 1b, while 1,25(OH)₂D₃ strongly induced the OC promoter-driven luciferase reporter, introduction of the expression vector for Grip NR boxes inhibited this transcriptional induction in a dose-dependent manner by as much as

a



b




Fig. 1. Gal4 DBD-Src-1 and Gal4 DBD-Grip fusion proteins suppress 1,25(OH)₂D₃ stimulated transcription from the osteocalcin promoter in MC3T3 E1 cells. **a:** Peptide expression constructs employed. Each construct encodes an amino terminal Gal4 DNA binding domain (residues 1–147) fused to either Src-1 (residues 621–765 containing three LXXLL (NR) boxes) or Grip (residues 629–760 containing three LXXLL (NR) boxes). **b:** A total of 500 ng of DNA/well comprised of 250 ng of pOC-3900, 50 ng RSV-βGal, and up to 200 ng of pM-Grip (adjusted with pM alone) was transfected into MC3T3 E1 cells plated 24 h earlier. Cells were treated with vehicle or 1,25(OH)₂D₃ (10⁻⁸ M) for 24 h. Activities are the means of triplicate assays ± SD. b is significantly different from a, c, and d; *P* < 0.05.


60%. Since these peptides interact directly with the AF-2 domain of the nuclear receptor family [Hall et al., 2000], our results indicate that blockade of this region prevents transcriptional activation by the VDR (or RXR) and supports the prevalent idea that the receptor must recruit an additional factor(s) through that site to manifest function. They also establish a system, whereby LXXLL-containing peptides can be evaluated for antagonist properties.

LXXLL-Gal4 DBD Fusion Peptides Interact With VDR

Phage display screening was used previously with ERα and ERβ as nuclear receptor targets to identify short LXXLL containing peptides that

were capable of selectively blocking the transactivating ability of ERα or ERβ when fused to the DBD of Gal4 [Hall et al., 2000]. Many of these peptides interacted with additional nuclear receptors as well [Hall et al., 2000]. We selected a subset of these peptide-Gal4 DBD fusion genes whose sequences have been previously reported [Chang et al., 1999; Hall et al., 2000] and assessed their capacity to produce proteins capable of interacting with either VDR or RXRα in a mammalian two-hybrid system. As is evident in Figure 2a (left panel), both the Src-1 and Grip NR boxes as well as a number of peptide-Gal4 DBD fusion proteins were able to interact with the VDR in the presence of 1,25(OH)₂D₃ as indicated by levels of luciferase activity that are well above those observed with the Gal4 DBD control alone (pM). Positively interacting peptides, including C33, D47, and EBIP41 and EBIP44 (Fig. 2a, left panel) each varied in relative activity. Several peptides did

a



b




Fig. 2. Gal4 DBD LXXLL fusion proteins interact with VDR and/or RXRα in a mammalian two-hybrid system. A total of 600 ng of DNA/well comprised of 250 ng of pGal4(5x)luc3, 50 ng RSV-βGal, 50 ng of either pM vector or the indicated fusion proteins and 50 ng of either pHVDR-VP16 (a) or pRXRα-VP16 (b) was transfected into MC3T3 E1 cells plated 24 h earlier. Cells were treated without or with 1,25(OH)₂D₃ (10⁻⁸ M) (A) or 9-*cis* RA (10⁻⁶ M) (B) as indicated, harvested 24 h later and evaluated for both β-galactosidase and luciferase activities. Activities are the means of triplicate assays ± SD. A: b is different from a, *P* < 0.05. B: Left panel: b is significantly different from a, *P* < 0.05; right panel: b, c, and d are significantly different from a, *P* < 0.05 and c is significantly different from d, *P* < 0.05.

not interact with the VDR including EBIP45, EBIP66, EBIP92, and F6 (Fig. 2a (left panel). The activities of several of the chimeric peptides in the absence and presence of $1,25(\text{OH})_2\text{D}_3$ are observed in Figure 2a (right panel). In the absence of ligand, the basal activity for these peptides as well as those seen in the left panel of Figure 2a was similar to that seen with the Gal 4 DBD alone (pM vector).

LXXLL-Gal4 DBD Fusion Peptides That Interact With RXR α

Examination of each of these peptides in the presence of RXR α -VP16 and 9-*cis* RA revealed that while several of the VDR reactive peptides also bound to RXR (C33, D47, EBIP41, and EBIP44 (Fig. 2b, left panel)), a number of peptides that were unreactive to VDR interacted significantly with RXR α including F6 and EBIP66 (Fig. 2b, left panel). As with VDR, EBIP92 showed no reactivity with RXR. Interestingly, while RXR α exhibited only basal activity with most of the peptides in the absence of 9-*cis* RA, its interaction with the F6 peptide was significantly above baseline (Fig. 2b, right panel). This suggests that the Gal 4 DBD-F6 protein can interact with unliganded and liganded RXR α . The results of these studies with VDR and RXR α indicate that the peptides examined exhibit three patterns of reactivity: interaction with (1) both RXR α and VDR, (2) RXR α alone, and (3) neither receptor.

Inhibition of $1,25(\text{OH})_2\text{D}_3$ or 9-*cis* RA Response With LXXLL-Gal4 DBD Chimeric Proteins

Having identified several LXXLL peptides that appear to exhibit VDR and RXR α interaction selectivity, we then assessed whether these peptides could block either VDR- or RXR α -induced transactivation. We used the human OC promoter (as in Fig. 1) as a target for VDR, whereas we used the TK promoter containing three copies of the RXRE derived from the CRABP II gene [Mangelsdorf et al., 1991] as a target for RXR. RXR expression in MC3T3 E1 cells was insufficient to mediate 9-*cis* RA response such that endogenous levels were supplemented through the use of an RXR α expression vector. As seen in Figure 3, while $1,25(\text{OH})_2\text{D}_3$ -stimulated the OC promoter strongly in the absence of peptide (pM vector alone), the VDR-reactive LXXLL peptides Src-1, Grip, C33, D47, and EBIP41, and EBIP44 all suppressed the




Fig. 3. Gal4 DBD fusion peptides C33, D42, D47 but not E92, suppress $1,25(\text{OH})_2\text{D}_3$ -stimulated transcription from the osteocalcin promoter in MC3T3 E1 cells. A total of 500 ng of DNA/well comprised of 250 ng of pOC-3900, 50 ng RSV- β Gal, and 200 ng of either pM-Src-1, pM-Grip, pM-C33, pM-D47, pM-EBIP41, pM-EBIP44, or pM-EBIP92 (adjusted with pM alone) was transfected MC3T3 E1 cells plated 24 h earlier. Cells were treated with vehicle or $1,25(\text{OH})_2\text{D}_3$ (10^{-8} M) for 24 h, harvested and evaluated for both β -galactosidase and luciferase activities. Activities are the means of triplicate assays \pm SD. b is different from a, $P < 0.05$.

receptor's activity when used at a concentration of 200 ng of plasmid. The findings in Figure 4 reveal that peptide Src-1 as well as the RXR α -reactive peptides F6 and EBIP66 each suppressed 9-*cis* RA-induced activity compared to the unreactive control peptide EBIP92. These results indicate that single copies of receptor-specific LXXLL peptides are capable of interacting with nuclear receptor targets with sufficient affinity to antagonize normal transcriptional ability.

RXR α -Selective LXXLL Peptides Block VDR-Mediated Transactivation

RXR is believed to participate directly as a heterodimeric partner in VDR mediated transactivation [Sone et al., 1991a; Yu et al., 1991; Mangelsdorf et al., 1995; Whitfield et al., 1995; Jin et al., 1996]. Thus, a peptide capable of inhibiting 9-*cis* RA-activated, RXR-mediated




Fig. 5. Gal4 DBD fusion peptides F6 and E66 suppress 1,25(OH) $_2$ D $_3$ -stimulated transcription from the osteocalcin promoter in MC3T3 E1 cells. A total of 500 ng of DNA/well comprised of 250 ng of pOC-3900, 50 ng RSV- β Gal, and up to 200 ng of either pM-Grip, pM-E92, pM-E53, pM-E60, pM-E66, or pM-F6 (adjusted with pM to total 600 ng) was transfected into MC3T3 E1 cells plated 24 h earlier. Cells were treated with vehicle or 1,25(OH) $_2$ D $_3$ (10^{-8} M) for 24 h, harvested and evaluated for both β -galactosidase and luciferase activities. Activities are the means of triplicate assays \pm SD. b is significantly different from a, $P < 0.05$.

several LXXLL motifs isolated from the Src1 and Grip comodulators as well as from a phage display screen can interact with the AF2 domain of the VDR and can function as antagonists when linked to a chimeric protein such as the Gal 4 DBD. We also demonstrate that LXXLL peptides directed specifically towards RXR and capable of suppressing 9-*cis* RA stimulated transcription also inhibit 1,25(OH)₂D₃ response. Our observations support the current hypothesis that the AF2 domain of the VDR is essential for 1,25(OH)₂D₃ stimulated transactivation and that RXR not only functions as a heterodimeric partner of VDR but may also facilitate transactivation.

RXR is widely believed to be a necessary partner for VDR activity in response to 1,25(OH)₂D₃ [Sone et al., 1991a,b; Whitfield et al., 1995; Cheskis and Freedman, 1996; Jin et al., 1996; Lemon and Freedman, 1996; Thompson et al., 1998; Thompson et al., 2001]. Despite this evidence, the absolute requirement for RXR in diverse 1,25(OH)₂D₃-stimulated responses has not been determined largely because the RXRs are derived from three different genes and

The VDR functions in response to $1,25(\text{OH})_2\text{D}_3$ to alter the expression of specific genes in a tissue-selective fashion within cells [Evans, 1989; Haussler et al., 1998]. The nuclear receptor family, of which the VDR is a member, acts to recruit additional comodulators to regulated promoters [Lemon and Freedman, 1996]. This recruitment involves an activation domain on the receptor surface and an LXXLL motif located within the comodulator itself [McInerney et al., 1998; Hall et al., 2000]. We show here that

almost ubiquitously expressed in mammalian cells [Evans, 1989; Mangelsdorf et al., 1995]. Our current studies lend additional support for the above prevailing idea by showing that inhibition of endogenous RXR in cells (EBIP66 and F6) can limit vitamin D response. Several of the RXR-selective peptides (EBIP53 and EBIP60) did not inhibit vitamin D response. It is possible that these particular peptides do not display an affinity for RXR that is sufficient to prevent recruitment of endogenous coactivators and therefore cannot block vitamin D response.

RXR is also believed to participate directly in the transactivation process [Thompson et al., 2001]. Accordingly, mutations in the activation domain of RXR were found to blunt the ability of VDR to activate transcription, thus acting as dominant negative inhibitors of VDR action. These findings indicate that both receptors likely participate in comodulator interaction essential to transactivation. Our studies support this view in that LXXLL peptides can both interact with the RXR AF-2 and block VDR transcriptional activity. Interestingly, the residues within the RXR AF2 region that mediate RXR homodimer and RXR/VDR heterodimer activity are different. This may reflect a difference in the organization of the receptors on unique DNA sequence elements, promoter context or, alternatively, selectivity in the cofactors that are recruited to those surfaces. The ability of peptides to exhibit selectivity between VDR and RXR, as seen in these studies, may also be indicative of receptor preferences for endogenous comodulators.

Ligands activate nuclear receptors by inducing receptor conformations capable of subsequent interaction with LXXLL containing coactivators. In view of this, while the F6 peptide can be argued to interact with both unliganded and liganded RXR, it is surprising that peptides that interact in a 9-*cis* RA-dependent manner with RXR such as EBIP66 can block a vitamin D-induced response in the absence of the RXR ligand. This finding suggests the possibility that VDR itself may act to induce a conformation within RXR that is similar although perhaps not identical to that induced by 9-*cis* RA. This observation supports a similar conclusion drawn from the finding that specific mutations within the LXXLL binding site of RXR (different from those that inhibit 9-*cis* RA dependent RXR activity) block vitamin D response [Thompson et al., 2001].

Vitamin D exerts highly pleiotropic actions on a variety of cell types through the regulation of numerous genes [Haussler et al., 1998; DeLuca and Cantorna, 2001]. While the actions of VDR as an RXR heterodimer on genes such as OP, OC, and 24OHase appear well supported, it is possible that additional regulatory paradigms may exist where the VDR functions differently. For example, recent evidence suggests that the VDR may function in certain settings as a monomer. Accordingly, the VDR appears to bind to an unusual DNA sequence half site on the GM-CSF promoter and to interact with proteins other than RXR such as NFAT1 and c-Jun [Towers and Freedman, 1998]. In addition, VDRE sequences that mediate suppression in the human parathyroid hormone gene may also require a protein other than RXR [Mackey et al., 1996]. These studies support the idea that the VDR could function through several different mechanisms to regulate transcription. Thus, RXR-selective peptides that are capable of inhibiting responses that require RXR such as those identified herein may be useful in defining the role of RXR in a wide variety of individual vitamin D-induced activities. Current efforts focus on using these peptides to inhibit specific endogenous responses to 1,25(OH)₂D₃ in cells and in vivo.

ACKNOWLEDGMENTS

This work was supported by DK-52453 (JWP) from the National Institutes of Health, CA 90645 (DPM) from the National Cancer Institute and DAMD17-99-1-9173 (C-YC) from the Department of Defense.

REFERENCES

- Baker AR, McDonnell DP, Hughes M, Crisp M, Mangelsdorf DJ, Haussler MR, Shine J, Pike JW, O'Malley BW. 1988. Molecular cloning and expression of human vitamin D₃ receptor complementary DNA: Structural homology with thyroid hormone receptor. *Proc Natl Acad Sci USA* 85:3294–3298.
- Brzozowski AM, Pike AC, Dauter Z, Hubbard RE, Bonn T, Engstrom O, Ohman L, Greene GL, Gustafsson JA, Carlquist M. 1997. Molecular basis of agonism and antagonism in the oestrogen receptor. *Nature* 389:753–758.
- Burmester JK, Wiese RJ, Maeda N, DeLuca HF. 1988. Structure and regulation of the rat 1,25-dihydroxyvitamin D₃ receptor. *Proc Natl Acad Sci USA* 85:9499–9502.
- Chang C, Norris JD, Gron H, Paige LA, Hamilton PT, Kenan DJ, Fowlkes D, McDonnell DP. 1999. Dissection of the LXXLL nuclear receptor-coactivator interaction motif using combinatorial peptide libraries: Discovery

- of peptide antagonists of estrogen receptors alpha and beta. *Mol Cell Biol* 19:8226–8239.
- Cheskis B, Freedman LP. 1996. Modulation of nuclear receptor interactions by ligands: Kinetic analysis using surface plasmon resonance. *Biochemistry* 35:3309–3318.
- DeLuca HF, Cantorna MT. 2001. Vitamin D: Its role and uses in immunology. *FASEB J* 15:2579–2585.
- Evans RM. 1989. The steroid and thyroid hormone receptor superfamily. *Science* 240:889–895.
- Gampe RT, Jr., Montana VG, Lambert MH, Miller AB, Bledsoe RK, Milburn MV, Kliewer SA, Willson TM, Xu HE. 2000. Asymmetry in the PPARgamma/RXRalpha crystal structure reveals the molecular basis of heterodimerization among nuclear receptors. *Mol Cell* 5:545–555.
- Hall JM, Chang CY, McDonnell DP. 2000. Development of peptide antagonists that target estrogen receptor beta-coactivator interactions. *Mol Endocrinol* 14:2010–2023.
- Haussler MR, Whitfield GK, Haussler CA, Hsieh J-C, Thompson PD, Selznick SH, Encinas Dominguez C, Jurutka PW. 1998. The nuclear vitamin D receptor: Biological and molecular regulatory properties revealed. *J Bone Min Res* 13:325–349.
- Jin CH, Kerner SA, Hong MH, Pike JW. 1996. Transcriptional activation and dimerization functions in the human vitamin D receptor. *Mol Endocrinol* 10:945–957.
- Kerner SA, Scott RA, Pike JW. 1989. Sequence elements in the human osteocalcin gene confer basal activation and inducible response to hormonal vitamin D₃. *Proc Natl Acad Sci USA* 86:4455–4459.
- Lemon BD, Freedman LP. 1996. Selective effects of ligands on vitamin D₃ receptor- and retinoid X receptor-mediated gene activation in vivo. *Mol Cell Biol* 16:1006–1016.
- Love JD, Gooch JT, Benko S, Li C, Nagy L, Chatterjee VKK, Evans RM, Schwabe JWR. 2002. The structural basis for the specificity of retinoid-X-receptor selective agonists: New insights into the role of helix H12. *J Biol Chem* 277:11385–11391.
- Mackey SL, Heymont JL, Kronenberg HM, Demay MB. 1996. Vitamin D receptor binding to the negative human parathyroid hormone vitamin D response element does not require the retinoid X receptor. *Mol Endocrinol* 10:298–305.
- Malloy PF, Pike JW, Feldman D. 1999. The vitamin D receptor and its role in hereditary 1,25-dihydroxyvitamin D₃ resistant rickets. *Endocr Rev* 20:156–188.
- Mangelsdorf DJ, Umersono K, Kliewer SA, Borgmeyer U, Ong ES, Evans RM. 1991. A direct repeat in the cellular retinol-binding protein type II gene confers differential regulation by RXR and RAR. *Cell* 66:555–561.
- Mangelsdorf DJ, Thummel C, Beato M, Herrlich P, Schutz G, Umersono K, Blumberg B, Kastner P, Mark M, Chambon P, Evans RM. 1995. The nuclear receptor superfamily: The second decade. *Cell* 83:835–839.
- McInerney EM, Rose DW, Flynn SE, Westin S, Mullen TM, Kronen A, Inostroza J, Torchia J, Nolte RT, Assa-Munt N, Milburn MV, Glass CK, Rosenfeld MG. 1998. Determinants of coactivator LXXLL motif specificity in nuclear receptor transcriptional activation. *Genes Dev* 12:3357–3368.
- Noda M, Vogel RL, Craig AM, Prah J, DeLuca HF, Denhardt DT. 1990. Identification of a DNA sequence responsible for binding of the 1,25-dihydroxyvitamin D₃ receptor and 1,25-dihydroxyvitamin D₃-enhancement of mouse secreted phosphoprotein 1 (Spp-1 or osteopontin) gene expression. *Proc Natl Acad Sci USA* 87:9995–9999.
- Perissi V, Staszewski LM, McInerney EM, Kurokawa R, Kronen A, Rose DW, Lambert MH, Milburn MV, Glass CK, Rosenfeld MG. 1999. Molecular determinants of nuclear receptor-corepressor interaction. *Genes Dev* 13:3198–3208.
- Rachez C, Freedman LP. 2000. Mechanisms of gene regulation by vitamin D₃ receptor: A network of coactivator interactions. *Gene* 246:9–21.
- Rachez C, Lemon BD, Suldan Z, Bromleigh V, Gamble M, Naar AM, Erdjument-Bromage H, Tempst P, Freedman LP. 1999. Ligand-dependent transcription activation by nuclear receptors requires the DRIP complex. *Nature* 398:824–828.
- Rachez C, Gamble M, Chang CP, Atkins GB, Lazar MA, Freedman LP. 2000. The DRIP complex and SRC-1/p160 coactivators share similar nuclear receptor binding determinants but constitute functionally distinct complexes. *Mol Cell Biol* 20:2718–2726.
- Ryu S, Zhou S, Ladurner AG, Tjian R. 1999. The transcriptional cofactor complex CRSP is required for activity of the enhancer-binding protein Sp1. *Nature* 397:446–450.
- Shiau AK, Barstad D, Loria PM, Cheng L, Kushner PJ, Agard DA, Greene GL. 1998. The structural basis of estrogen receptor/coactivator recognition and the antagonism of this interaction by tamoxifen. *Cell* 95:927–937.
- Sone T, Kerner SA, Pike JW. 1991a. Vitamin D receptor interaction with specific DNA: Association as a 1,25-dihydroxyvitamin D₃-modulated heterodimer. *J Biol Chem* 266:23296–23305.
- Sone T, Ozono K, Pike JW. 1991b. A 55 kilodalton accessory factor facilitates vitamin D receptor DNA binding. *Mol Endocrinol* 5:1578–1586.
- Thompson PD, Jurutka PW, Haussler CA, Whitfield GK, Haussler MR. 1998. Heterodimeric DNA binding by the vitamin D receptor and retinoid X receptors is enhanced by 1,25-dihydroxyvitamin D₃ and inhibited by 9-cis-retinoic acid. Evidence for allosteric receptor interactions. *J Biol Chem* 273:8483–8491.
- Thompson PD, Remus LS, Hsieh JC, Jurutka PW, Whitfield GK, Galligan MA, Encinas Dominguez C, Haussler CA, Haussler MR. 2001. Distinct retinoid X receptor activation function-2 residues mediate transactivation in homodimeric and vitamin D receptor heterodimeric contexts. *J Mol Endocrinol* 27:211–227.
- Towers TL, Freedman LP. 1998. Granulocyte-macrophage colony-stimulating factor gene transcription is directly repressed by the vitamin D₃ receptor. Implications for allosteric influences on nuclear receptor structure and function by a DNA element. *J Biol Chem* 273:10338–10348.
- Whitfield GK, Hsieh J-C, Nakajima S, MacDonald PN, Thompson PD, Jurutka PW, Haussler CA, Haussler MR. 1995. A highly conserved region in the hormone binding domain of the human vitamin D receptor contains residues vital for heterodimerization with retinoid X receptor and for transcriptional activation. *Mol Endocrinol* 9:1166–1179.
- Yu V, Delsert C, Anderson B, Holloway JM, Devary OV, Naar AM, Kim SY, Boutin JM, Glass CK, Rosenfeld MG. 1991. RXRβ: A coregulator that enhances binding of retinoic acid, thyroid hormone, and vitamin D receptors to their cognate response elements. *Cell* 67:1251–1266.